arXiv:cond-mat/9912156v1 [cond-mat.mes-hall] 9 Dec 1999

On the response of composite fermions to weak electrostatic potentials

Behnam Farid
Cavendish Laboratory, Department of Physics, University of Cambridge,
Madingley Road, Cambridge CB3 0HE, United Kingdom
(February 1, 2008)

We establish that the response to static perturbations of two-dimensional electron systems (2DESs), exposed to magnetic field $\boldsymbol{B}$, in states with the Landau-level filling fractions $\nu_{\rm e}$ close to one such as 1/2, is singularly different from that of 2DESs at small $\|\boldsymbol{B}\|$. Thus in addition to demonstrating the sever inadequacy of the treatment of composite fermions as non-interacting particles, we show, in contradiction to a recent theoretical finding, that the physical origin of the observed behaviour of the dc longitudinal magneto-resistivity for $\nu_{\rm e} \approx 1/2$ in periodically modulated 2DESs remains to be determined.

71.10.Pm, 73.40.Hm, 73.50.Jt

Advance towards understanding of the fractional quantum Hall effect (FQHE) was made through the variational wavefunction due to Laughlin [1] for the corresponding states. Subsequent progress was made through the application of a statistical transformation [2], transmuting electrons into hard-core bosons, by Girvin and MacDonald [3] who showed that the equal-time boson-boson correlation function of the transformed Laughlin wavefunction exhibits algebraic off-diagonal long-range order. This statistical transformation has been an essential ingredient in considerations by the latter authors [3], Read [4] and Zhang, Hansson and Kivelson [5] in formulating a Ginzburg-Landau theory for the FQHE. The microscopic formulation of this theory relies on a statistical gauge field associated with the Chern-Simons (CS) action in $2+1$ dimensions whose coupling constant $\Theta_{\rm CS}$ is equal to an odd multiple of 2π.

An alternative approach to the FQHE was advanced by Jain [6], who through introducing a variational wavefunction, involving the Laughlin wavefunction pertaining to a filled Landau level (LL) to whose constituent particles an *even* number of flux quanta are attached, ascribed the phenomenon of FQHE to the integer QHE (IQHE) of 'composite fermions' (CFs). A CF is thus composed of an electron and $2n$, $n = 1, 2, \ldots$, quantised magnetic flux tubes of magnetic flux $\Phi_0 \equiv h/e$ (throughout, $-e < 0$). The total magnetic flux due to these flux tubes results in a reduced effective magnetic flux density $\Delta\boldsymbol{B}$ of whose corresponding LLs (for CFs) in the case where the *electronic* $\nu_{\rm e}$ is equal to $p/(2np+1)$, exactly p levels are completely filled *and* none is fractionally occupied. We note in passing that for $\nu_{\rm e} \in (0, 1]$, the 'particle-hole' symmetry (which is an exact property for the case where the lowest LL is decoupled from the higher LLs) implies that $\nu_{\rm e}' {:=} 1 - \nu_{\rm e}$ corresponds to a FQH state if $\nu_{\rm e}$ does to one, so that the Jain series $\nu_{\rm e} = p/(2np+1)$ implies FQH states at $\nu_{\rm e}' = 1 - p/(2np+1)$; each of these, on account of the 'law of corresponding states' [7], in turn implies FQH states at $\nu_{\rm e}'' = m + \nu_{\rm e}$, where m is an integer. For increasing values of p, $\nu_{\rm e} = p/(2np+1)$ and $\nu_{\rm e}' {:=} 1 - \nu_{\rm e}$ approach $\nu_{\rm e} = 1/(2n)$ from left and right, respectively. The filling fraction $\nu_{\rm e} = 1/(2n)$, although a limit of those associated with FQH states for $p \to \infty$, does *not* itself correspond to a QH state, in that its associated state is compressible [8]. The notable fractions in this category are $\nu_{\rm e} = 1/2, 1/4$ and $\nu_{\rm e}' = 3/4$.

The scenario as put forward by Jain [6] finds its microscopic justification in the fermionic version of the CS field theory in $2+1$ dimensions where $\Theta_{\rm CS} = 2n \times 2\pi$, $n = 1, 2, \ldots$. Indeed, from the mean-field Hamiltonian of this theory for *extended* 2DESs [9], [10], [8], it is apparent that CFs are exposed to an effective magnetic flux density equal to ΔB (here and further on, B and ΔB denote the amplitudes of $\boldsymbol{B}$ and $\Delta\boldsymbol{B}$ which are assumed to be perpendicular to the plane of 2DESs). A most appealing aspect of the fermionic CS field theory in $2+1$ dimensions was exposed by Lopez and Fradkin [11] who demonstrated that all physical amplitudes calculated within this theory are identical with those calculated within the conventional theory where $\Theta_{\rm CS} = 0$. It follows that in cases where $\Theta_{\rm CS} = 2n \times 2\pi$, the CS field theory has the *practical* advantage that its mean-field approximation captures the essential aspects of the FQH states, such as their incompressibility. CFs have found experimental support in the surface-acoustic-wave resonance experiments by Willett, *et al.* [12], geometric resonance measurements by Kang, *et al.* [13] and Willett,*et al.* [14] and magnetic focusing experiments by Goldman, *et al.* [15] and Smet, *et al.* [16].

A recent experimental work by Smet, *et al.* [17] on periodically and supposedly *weakly* modulated 2DESs has revealed some unexpected aspects concerning the dc longitudinal magneto-resistivity ϱ_{xx} as function of B close to $\nu_{\rm e} = 1/2$. These include the observation that a supposedly 1% modulation amplitude in the number density n_e of electrons gave rise to a positive magneto-resistance, for $|B - B_{1/2}|$, typical of a charge-density modulation amplitude of 10% (see below, in particular the paragraph containing the definition for ϵ); here $B_{\nu_{\rm e}} \equiv \Phi_0 n_e/\nu_{\rm e}$ is the magnetic flux density corresponding to $\nu_{\rm e}$. In this connection, the authors [17] state "Clearly some ingredients are missing to form a consistent model, unless our implicit assumption, that the strength of the density perturbation taken from low field data can be transferred to the high field regime, is incorrect." One of the findings of the present work is that this is indeed the case.

The observations by Smet, *et al.* [17], have been the subject of theoretical consideration by von Oppen, *et al.* [18], Mirlin, *et al.* [19], and Zwerschke and Gerhardts [20]. By taking into account the differences in the scattering properties of random electrostatic and magnetic fields (in the case at hand the two are related via the relationship $\Delta B \equiv \Phi_0 n_e/p$), Mirlin, *et al.* [19] and Zwerschke and Gerhardts [20] obtain results for ϱ_{xx}, concerning a periodically modulated 2DES along the x direction, which are in reasonable agreement with experimental results. By further solving a linearised Boltzmann equation for CFs which in a self-consistent manner takes account of the electric field generated by the motion of the flux tubes associated with CFs, the latter authors obtain results which are in excellent agreement with experiment. A crucial observation by Zwerschke and Gerhardts [20] is that the so-called 'channeled orbits', which are neglected in the approximate treatment of the Boltzmann equation due to Beenakker [21], determine the specific v-shape in the experimentally observed $\varrho_{xx}(B)$ close to $B = B_{1/2}$.

Since in the above-cited works the CFs are considered as non-interacting particles, the strength of the modulating potential as 'seen' by CFs is not the bare one, namely $v_{\rm ext}(x)$, but one which is screened, namely $v_{\rm eff}(x)$. This becomes evident by applying the mean-field approximation subsequent to having introduced $v_{\rm ext}(x)$ into the many-body Hamiltonian of an otherwise uniform system. Assuming $v_{\rm ext}(x) = v_Q \cos(Qx)$, with $Q = 2\pi/a_x$, *linear* screening suffices for small v_Q and one has $v_{\rm eff}(x) = \big(\varepsilon_{\rm pp}^{-1}(Q,0) v_Q\big) \cos(Qx)$, where $\varepsilon_{\rm pp}^{-1}(Q,0)$ stands for the inverse of the static 'proton-proton' dielectric function [22]; $\varepsilon_{\rm pp}^{-1}(q,0) = 1 + \big(v(q) + \kappa_{\rm xc}(q)\big)\chi(q,0)$ where χ denotes the density-density response function of the interacting system and $v(q)$ the Fourier transform of the electron-electron interaction function which in this work we assume to be the Coulomb interaction function so that $v(q) = e^2/(2\epsilon_0\epsilon_r q)$ with ϵ_0 the vacuum permittivity and ϵ_r for the relative dielectric constant; $\kappa_{\rm xc}(q)$ is the exchange and correlation contribution to the interaction. For the ε^{-1} specific to screening of the electron-electron interaction, one has $\varepsilon^{-1}(q,0) = 1 + v(q)\chi(q,0)$. Since $\kappa_{\rm xc}(q)/v(q) = \mathcal{O}(q)$ for $q \to 0$ and since in our considerations $Q \ll k_F$ (we consider $n_e = 1.36 \times 10^{11}$ cm^{-2} and $a_x = 400$ nm), with $\boldsymbol{k}_F$ the Fermi wavevector, we can neglect $\kappa_{\rm xc}(q)$ and thus identify $\varepsilon_{\rm pp}^{-1}$ with ε^{-1}.

A modulation is considered weak if the dimensionless quantity $\epsilon := e\varepsilon_{\rm pp}^{-1}(Q,0) v_Q/(\sqrt{2}E_F)$, is small ($\epsilon$ as defined here is identical with that by Beenakker [21] and is $1/\sqrt{2}$ times the ϵ by Zwerschke and Gerhardts [20] which coincides with η by Mirlin, *et al.* [19]). This is because according to the Thomas-Fermi approximation (TFA), $\varepsilon_{\rm TF}^{-1}(q,0) = 1/(1 + g_e(E_F)v(q))$ where $g_e(\omega) = m_b/(\pi\hbar^2)$ is the *total* density of one-electron states for spin-unpolarised 2DES, with m_b the bare band-electron mass; with $v(q)$ the Coulomb interaction, $g_e(E_F)v(q) = a_0 q/2$, where $a_0 := 4\pi\epsilon_0\epsilon_r\hbar^2/(m_b e^2)$ is the effective Bohr radius (in this work, $a_0 = 9.79 \times 10^{-9}$ m), so that with $a_x \approx 400 \times 10^{-9}$ m, $\varepsilon_{\rm TF}^{-1}(Q,0) \approx a_0 Q/2 \ll 1$ *and* $\chi_{\rm TF}(Q,0) \approx -1/v(Q)$ from which, making use of $E_F g_e = n_e$, one obtains $\delta n_e(Q)/n_e \equiv e\chi(Q,0)v_Q/n_e \approx \sqrt{2}\epsilon$. By *assuming* the validity of the TFA for CFs, one has $\epsilon_{\rm CF} \approx \epsilon$ (with $m_\star \sim 5m_b$ the CF effective mass, $\varepsilon_{\rm TF}^{-1}(Q,0) \approx (2m_b/m_\star)a_0 Q/2 \ll 1$, so that for CFs also $\chi_{\rm TF}(Q,0) \approx -1/v(Q)$). Thus the *assumption* with regard to the validity of the TFA for CFs implies that a 'weak' perturbation for electrons is equally 'weak' for CFs and vice versa. It is common practice to assign a fixed, i.e. independent of B, value to ϵ, reflecting the constancy of ϵ within the TFA. The significance of ϵ is clarified by considering the expression for ϱ_{xx} as determined by Beenakker (Eq. (6) in Ref. [21]); by choosing $\epsilon = 0.015$ [21], this ϱ_{xx} accurately describes the magneto-resistance oscillations as observed by Weiss, *et al.* [23].

For a 2DES in the paramagnetic phase exposed to magnetic field, we have $\chi(q,\omega) = \chi^0(q,\omega)/\big(1 - (v(q) + \kappa_{\rm xc}(q))\chi^0(q,\omega)\big)$, where [24]

$$\chi^0(q,0) = \frac{-m_b}{\hbar^2\pi^2}\mathcal{F}_{[E_F/(\hbar\omega_c)]}\left(\frac{\hbar q^2}{2m_b\omega_c}\right); \qquad (1)$$

here $[x]$ denotes the greatest integer less than or equal to x and $\mathcal{F}_N(a) := \int_0^\pi {\rm d}\tau\ \tau \sin(a\sin\tau)\exp(-a[1 + \cos\tau])G_N(2a[1 + \cos\tau])$; $G_N(x) \equiv 1$, when $N = 0$, and $G_N(x) \equiv L_N(x) + 2L_{N-1}^{(1)}(x)$, when $N \geq 1$. Here $L_N(x)$ denotes the Nth-order Laguerre polynomial and $L_N^{(m)}(x) := (-1)^m {\rm d}^m L_{N+m}(x)/{\rm d}x^m$. By neglecting $\kappa_{\rm xc}(q)$, the expression for $\chi(q,0)$ reduces to the random-phase approximation (RPA). Since in our applications in the present work $Q \ll k_F$, we neglect $\kappa_{\rm xc}$ and thus employ the RPA for $\chi(q,0)$.

For CFs, the counterpart of $\chi(q,\omega)$ is denoted by $-K_{00}(q,\omega)$ which is the $(\mu,\nu) = (0,0)$ component of a 2×2 matrix $\mathbf{K}(q,\omega)$ with $K_{\mu,\nu} \equiv (\mathbf{K})_{\mu,\nu}$. Here $\mu,\nu \in \{0,1\}$, with 0 indicating the 'time' component and 1 the transverse 'space' component [8]. Within the RPA, we have $\mathbf{K} = \mathbf{K}^0\big(\mathbf{I} + \mathbf{U}\mathbf{K}^0\big)^{-1}$, where $\mathbf{U} := \mathbf{V} + \mathbf{C}^{-1}$ in which $V_{00} = v(q)$ and $V_{\mu,\nu} \equiv 0$ for $(\mu,\nu) \neq (0,0)$, and for $\nu_{\rm e} = p/(2np+1)$, $C_{00} \equiv C_{11} \equiv 0$, $C_{01} \equiv -C_{10} \equiv iq/(h\tilde{\phi})$, where $\tilde{\phi} := 2n$ [8], [9], [25]. It can be shown [25] that $\mathbf{K}^0 = \mathbf{T}^{-1}\mathbf{S}\mathbf{T}^{-1}$, where $T_{00} \equiv i(e/q)\sqrt{i\omega}$, $T_{11} \equiv e/\sqrt{i\omega}$, $T_{01} \equiv T_{10} \equiv 0$ and $S_{00} \equiv ipe^2(\omega/\Delta\omega_c)\Sigma_0/h$, $S_{11} \equiv ipe^2(\Delta\omega_c/\omega)(\Sigma_2 + 1)/h$ and $S_{10} \equiv -S_{01} \equiv pe^2\Sigma_1/h$, in which $\Delta\omega_c := e\Delta B/m_b$; for $\omega = 0$ and $p \geq 1$,

$$\Sigma_j \equiv \frac{-\exp(-Y)}{p}\sum_{\ell=0}^{p-1}\sum_{m=p}^{\infty}$$

$$\{\frac{\ell!}{(m-\ell)m!}Y^{m-\ell-1}\{L_\ell^{(m-\ell)}(Y)\}^{2-j}$$

$$\times\{(m-\ell-Y)L_\ell^{(m-\ell)}(Y)+2Y\mathrm{d}L_\ell^{(m-\ell)}(Y)/\mathrm{d}Y\}^j\}, \quad (2)$$

where $Y{:=}(qR_\Delta)^2/(4p)$, with $R_\Delta \equiv \hbar k_F/(e\Delta B)$, the effective semi-classical cyclotron radius for CFs. Since the state of CFs corresponds to that of fully-polarised electrons, $k_F = \sqrt{4\pi n_e}$. One obtains

$$K_{00}(q,\omega=0) = \frac{-pq^2\Sigma_0}{h\Delta\omega_c\Upsilon}, \quad (3)$$

where $\Upsilon{:=}(1+p\tilde{\phi}\Sigma_1)^2 - pq^2v(q)\Sigma_0/(h\Delta\omega_c) - p^2\tilde{\phi}^2\Sigma_0(\Sigma_2+1)$. Assuming the CFs to have the mass $m_\star$, a direct change of m_b into $m_\star$, or of $\Delta\omega_c$ into $\Delta\omega_{c\star}{:=}e\Delta B/m_\star$, in the above expressions, would result in a $K_{00}(q,\omega)$ that violates a requirement implied by a theorem due to Kohn [26], [5], [8]. Simon and Halperin [25] have put forward modifications, denoted by MRPA ('modified RPA') and M^{2}RPA, through which any desired mass can be assigned to CFs without ill consequences of the kind. It turns out that M^{2}RPA is superior to MRPA, however they are identical in so far as $K_{00}(q,\omega)$ is concerned. We indicate the MRPA for $\mathbf{K}$ by $\widetilde{\mathbf{K}}$. With $\mathbf{K}_\star^0$ denoting the $\mathbf{K}^0$ in which $\Delta\omega_c$ is replaced by $\Delta\omega_{c\star}$, we have $\widetilde{\mathbf{K}} = \mathbf{K}_\star^0(\mathbf{I} + \{\mathbf{U}+\mathbf{F}\}\mathbf{K}_\star^0)^{-1}$, where $F_{00} \equiv (m_\star - m_b)\omega^2/(n_e q^2)$, $F_{11} \equiv -(m_\star - m_b)/n_e$, $F_{01} \equiv F_{10} \equiv 0$ [25]. A straightforward calculation yields

$$\widetilde{K}_{00}(q,0) = K_{\star 00}(q,0) - \frac{\mathcal{N}_1\mathcal{N}_2}{\mathcal{D}}, \quad (4)$$

$$\mathcal{N}_1 \equiv \Big(\frac{pq}{h\Upsilon_\star}\Big)^2\frac{m_\star - m_b}{n_e} \times\{(1+p\tilde{\phi}\Sigma_1)\Sigma_1 - p\tilde{\phi}\Sigma_0(\Sigma_2+1)\}, \quad (5)$$

$$\mathcal{N}_2 \equiv \frac{pq}{\Delta\omega_{c\star}}\Sigma_0\{\frac{qv(q)}{h}\Sigma_1 - \frac{\tilde{\phi}}{q}\Delta\omega_{c\star}(\Sigma_2+1)\} +\Sigma_1\{1 - \frac{pq^2v(q)}{h\Delta\omega_{c\star}}\Sigma_0 + p\tilde{\phi}\Sigma_1\}, \quad (6)$$

$$\mathcal{D} \equiv 1 + \frac{(m_\star - m_b)p}{hn_e\Upsilon_\star} \times\{pq\Sigma_1\{\frac{qv(q)}{h}\Sigma_1 - \frac{\tilde{\phi}}{q}\Delta\omega_{c\star}(\Sigma_2+1)\} +\Delta\omega_{c\star}(\Sigma_2+1)\{1 - \frac{pq^2v(q)}{h\Delta\omega_{c\star}}\Sigma_0 + p\tilde{\phi}\Sigma_1\}\}. \quad (7)$$

Above $\Upsilon_\star$ is the counterpart of Υ.

In our calculations concerning CFs, we employ $\nu_\mathrm{e} = hn_e/(eB)$ and subsequently evaluate p according to $p = [\nu_\mathrm{e}/(1-2n\nu_\mathrm{e})]$ if $\nu_\mathrm{e} < 1/(2n)$, and $p = [(1-\nu_\mathrm{e})/(1-2n(1-\nu_\mathrm{e}))]$, if $\nu_\mathrm{e} > 1/(2n)$ (in the present work, $n=1$); for $[x]$ see text following Eq. (1) above. This procedure introduces some error, that is in general $p/(2np+1)$, or $1-p/(2np+1)$ if $\nu_\mathrm{e} > 1/(2n)$, determined from the thus-obtained p is not exactly equal to $\nu_\mathrm{e} = hn_e/(eB)$. However, explicit calculation shows that for the range of B considered in this work, the relative error in ν_e (which oscillates around zero and whose magnitude increases with increasing $|\Delta B|$) never exceeds 0.06%.



FIG. 1. (colour) *Left panel*: $\varepsilon^{-1}(q,0)$ for CFs ((a), (b) and (c)) and electrons ((d) and (e)); (a) and (b) ((d)) correspond to $\Delta B(B) = 0.2$ T while (c) ((e)) corresponds to $\Delta B(B) = 0.4$ T. (a) and (c) are based on $\widetilde{\mathbf{K}}$ with $m_\star = 5\times m_b$, while for curve (b) $m_\star = m_b \equiv 0.067 m_e$, where m_e is the electron mass in vacuum; the coincidence of the tangents of curves (a) and (b) in the neighbourhood of $q=0$ indicates that $\widetilde{K}_{00}(q,0)$ is indeed in agreement with the requirement of the Kohn theorem. Here $a_0 = 9.79$ nm, corresponding to $\epsilon_r = 12.4$ and m_b. All results concern $n_e = 1.36\times 10^{11}$ cm^{-2} so that $k_F^{\text{elec.}} = 0.91/a_0$, $k_F^{\text{CF}} = 1.28/a_0$ ($\equiv \sqrt{2}k_F^{\text{elec.}}$). *Right panel*: $\varepsilon^{-1}(q,0)$ as a function of ΔB (for CFs with $m_\star = 5\times m_b$ — curve (a)) and B (for electrons — curve (b)) at $q = 0.15/a_0$; this q coincides with $Q = 2\pi/a_x$ corresponding to $a_x \approx 400$ nm. Curve (c) is 5× (b).

In Figs. 1 and 2 we present results of our numerical calculations. The sever inadequacy of the TFA in particular for CFs is apparent. One observes that over considerable range of values of ΔB, $\varepsilon^{-1}(q,0)$, and therefore ϵ defined above, pertaining to CFs is by one order of magnitude larger than that for electrons. This observation is in line with the statement by Smet, *et al.* [17] quoted above. One further observes that in the case of CFs, ϵ strongly depends on ΔB. The similarities between $-K_{\star 00}^0(q,0) \equiv pq^2\Sigma_0/(h\Delta\omega_{c\star})$ and $\chi^0(q,0)$, which describe the number-density response to the *total* (i.e. external plus induced) potential for systems of CFs and electrons, respectively (Fig. 2, right panel), and lack hereof in the corresponding $\varepsilon^{-1}(q,0)$, is indicative of the significance of Σ_1 and Σ_2 which have no counterparts in the case of electrons in weak magnetic fields. We note in passing that for ΔB and B approaching zero, the oscillations of $-K_{\star 00}^0(q,0)$ and $\chi^0(q,0)$ shift towards $q=0$, so much so that in the limits $\Delta B = 0$ and $B = 0$ all oscillations collapse at $q=0$ and $-K_{\star 00}^0(q\to 0,0)$ and $\chi^0(q\to 0,0)$ take on non-vanishing constant values as expected from compressible liquids. As is clearly seen from the left panels of Figs. 1 and 2, for decreasing B and

ΔB the functions $\varepsilon^{-1}(q,0)$ are increasingly 'pushed' towards lower values of q, signalling the emergence of the mentioned compressible states at $B = 0$ and $\Delta B = 0$. We point out that the step-like behaviour in $\varepsilon^{-1}(q,0)$ in the right panel of Fig. 1 is associated with the well-known cusps in the total energies as functions of magnetic flux density of respective systems, mathematically most clearly evident from $[E_F/(\hbar\omega_c)]$, which implies truncation, in the expression for $\chi^0(q,0)$ in Eq. (1); the *flatness* of the plateaus of $\varepsilon^{-1}(q,0)$ corresponding to CFs, on the other hand, is an artifact of the error associated with our restriction of ν_{e} to forms $p/(2np+1)$ and $1-p/(2np+1)$.



FIG. 2. (colour) *Left panel*: Similar to Fig. 1 but over a restricted range of values of $\varepsilon^{-1}(q,0)$ (the markings (a)-(e) coincide with those in Fig. 1). Curves (f) and (g) are the TFAs for $\varepsilon^{-1}(q,0)$ for electrons and CFs (using $m_\star = 5\times m_b$), respectively. *Right panel*: The static density-density response functions of *non*-interacting CFs, $-K^0_{\star\,00}(q,0)$ ((a) and (c), concerning $\Delta B = 0.2, 0.4$ T, respectively), and that of electrons, $\chi^0(q,0)$ ((d) and (e) — $B = 0.2, 0.4$ T).

In conclusion, by calculating the response functions of electrons and CFs, we have established that describing the transport properties of CFs in spatially modulated systems along the same lines as non-interacting electrons in weak magnetic fields is inadequate. The (excellent) agreement between the theoretical results by Mirlin, *et al.* [19] and Zwerschke and Gerhardts [20] with experiments are therefore to be attributed to the freedom in the choice of the parameter(s) of the collision operator in the Boltzmann equation dealt with by the authors. We are presently investigating the transport properties of CFs along the lines as put forward by Kim, *et al.* [27].

It is a pleasure for me to thank Professor Peter B. Littlewood for invaluable support and kind hospitality at Cavendish Laboratory. With appreciation I acknowledge stimulating discussions with Dr Jurgen H. Smet and Professor Klaus von Klitzing at the initial stages of this work.

---

[1] R. B. Laughlin, Phys. Rev. Lett. **50**, 1395 (1983).
[2] F. Wilczek, Phys. Rev. Lett. **49**, 957 (1982); D. Arovas, J. R. Schrieffer, and F. Wilczek, Phys. Rev. Lett. **53**, 722 (1984).
[3] S. M. Girvin, and A. H. MacDonald, Phys. Rev. Lett. **58**, 1252 (1987).
[4] N. Read, Phys. Rev. Lett. **62**, 86 (1989).
[5] S.-C. Zhang, T. H. Hansson, and S. Kivelson, Phys. Rev. Lett. **62**, 82 (1989). See also D.-H. Lee, and S.-C. Zhang, Phys. Rev. Lett. **66**, 1220 (1991).
[6] J. K. Jain, Phys. Rev. Lett. **63**, 199 (1989); Phys. Rev. B **40**, 8079 (1989); Adv. Phys. **41**, 105 (1992).
[7] S. Kivelson, D.-H. Lee, and S.-C. Zhang, Phys. Rev. B **46**, 2223 (1992).
[8] B. I. Halperin, P. A. Lee, and N. Read, Phys. Rev. B **47**, 7312 (1993).
[9] Q. Dai, J. L. Levy, A. L. Fetter, C. B. Hanna, and R. B. Laughlin, Phys. Rev. B **46**, 5642 (1992).
[10] Y.-H. Chen, F. Wilczek, E. Witten, and B. I. Halperin, Int. J. Mod. Phys. **3**, 1001 (1989).
[11] A. Lopez, and E. Fradkin, Phys. Rev. B **44**, 5246 (1991).
[12] R. L. Willett, R. R. Ruel, K. W. West, and L. N. Pfeiffer, Phys. Rev. Lett. **71**, 3846 (1993).
[13] W. Kang, H. L. Störmer, and L. N. Pfeiffer, Phys. Rev. Lett. **71**, 3850 (1993).
[14] R. L. Willett, K. W. West, and L. N. Pfeiffer, Phys. Rev. Lett. **83**, 2624 (1999).
[15] V. J. Goldman, B. Su, and J. K. Jain, Phys. Rev. Lett. **72**, 2065 (1994).
[16] J. H. Smet, *et al.*, Phys. Rev. Lett. **77**, 2272 (1996).
[17] J. H. Smet, K. von Klitzing, D. Weiss, and W. Wegscheider, Phys. Rev. Lett. **80**, 4538 (1998).
[18] F. von Oppen, A. Stern, and B. I. Halperin, Phys. Rev. Lett. **80**, 4494 (1998).
[19] A. D. Mirlin, P. Wölfle, Y. Levinson, and O. Entin-Wohlman, Phys. Rev. Lett. **81**, 1070 (1998).
[20] S. D. M. Zwerschke, and R. R. Gerhardts, Phys. Rev. Lett. **83**, 2616 (1999).
[21] C. W. J. Beenakker, Phys. Rev. Lett. **62**, 2020 (1989).
[22] V. Heine, and D. Weaire, in *Solid State Physics*, **24** (Academic, New York, 1970).
[23] D. Weiss, K. von Klitzing, K. Ploog, and G. Weimann, Europhys. Lett. **8**, 179 (1989).
[24] M. L. Glasser, Phys. Rev. B **28**, 4387 (1983).
[25] S. H. Simon, and B. I. Halperin, Phys. Rev. B **48**, 17368 (1993); S. H. Simon, J. Phys. C **8**, 10127 (1996).
[26] W. Kohn, Phys. Rev. **123**, 1242 (1961).
[27] Y. B. Kim, P. A. Lee, and X.-G. Wen, Phys. Rev. B **52**, 17275 (1995).